


## **Historic, Archive Document**

Do not assume content reflects current scientific knowledge, policies, or practices.



aSB608  
.P65S52  
1978  
Reserve

# HOW TO

## Identify and Control

## RHIZOSPHAERA NEEDLECAST



North Central Forest Experiment Station  
Forest Service

U.S. Department of Agriculture


MAY 3 - 1979

CATALOGING = PREP.

**Rhizosphaera** needlecast, caused by the fungus **Rhizosphaera kalkhoffii**, seriously damages blue spruce trees in nurseries and Christmas tree plantations in the north-central States. Although some trees are killed, the primary damage has been premature needlecast, rendering the tree unmerchantable.

## SYMPTOMS

Current-year needles are infected in May and June. Symptoms appear in late fall or the following spring, when fruiting bodies of the **Rhizosphaera** fungus appear in the stomata of infected needles. When magnified, these stomata appear as fuzzy black dots instead of the usual white color. The infected 2-year-old needles turn yellow in July and purplish-brown in late August. Most needles are cast in late summer, although some adhere overwinter and produce spores the following spring.



Fruiting bodies of **Rhizosphaera kalkhoffii** protruding from stomata appear as small black dots.



Infected needles eventually turn purplish-brown.



**Rhizosphaera** needlecast damage is most severe on lower portion of tree.



## SPREAD

Hyaline conidia (spores  $4 \times 8 \mu$  in size) are released from the fruiting bodies during wet weather in late spring. Rain-splash disseminates them to infect other needles. Ironically, spores may also be spread from tree to tree on the tools used to shear Christmas tree plantations, especially if shearing is done when foliage is wet. However, most infections in plantations result from infected planting stock.

## CONTROL

### Cultural

- Use healthy planting stock.
- Identify disease early to minimize losses.
- Discard seedlings that have fruiting bodies in the stomata.
- Shear trees in healthy plantations first.
- Sterilize tools by dipping in denatured alcohol for 3 minutes.
- Avoid shearing infected trees when foliage is wet.

### Chemical


Trees should be sprayed with Bordeaux mixture 8-8-100 early in June and again early in July. Chlorothalonil is also effective, and is registered for use in Minnesota and Wisconsin.

## CULTURING

Sometimes the only sure way to identify this disease is to culture the causal fungus. Using normal, sterile isolation techniques, here is how to do it:

- Dip infected needles for 1 to 2 seconds in a 1:10 hypochlorite solution (household bleach and water) to sterilize the surface.
- Using a fine needle, remove portions of the fruiting body containing spores from the infected stomata.
- Place the fruiting bodies onto potato dextrose agar in petri dishes and incubate at 68°F for 5 to 10 days.

The mycelial culture is at first light tan in color. This gradually darkens in about 10 days to a yellowish brown. Advancing hyphae remain close to the agar surface and soon begin to produce abundant spores that bud off from any point along the hyphal cells.



Culture of **Rhizosphaera**.

**DARROLL D. SKILLING**

Principal Plant Pathologist

Copies available from —

**North Central Forest  
Experiment Station  
Folwell Avenue  
St. Paul, MN 55108**

**NA State & Private  
Forestry  
370 Reed Road  
Broomall, PA 19008**

(For further information on **Rhizosphaera** needlecast see: Skilling, D.D., and T.H. Nicholls. 1974. **Rhizosphaera** needlecast. American Christmas Tree Journal 18:21-23.)



**REVISED 1978**